

RESEARCH PAPER



## Environmental determinants of previtamin D synthesis in the United Arab Emirates

Adam J. Buckley <sup>a</sup>, Zara Hannoun<sup>a</sup>, Nader Lessan<sup>a</sup>, Michael F. Holick <sup>b</sup>, and Maha T. Barakat<sup>a</sup>

<sup>a</sup>Imperial College London Diabetes Centre, Research Department, Abu Dhabi, United Arab Emirates; <sup>b</sup>Boston University Medical Center, Section of Endocrinology, Diabetes, and Nutrition, Department of Medicine, Boston, MA, USA

### ABSTRACT

Despite abundant sunshine throughout the year, vitamin D deficiency is endemic in the UAE. Solar radiation within the UVB range of the spectrum is required for the photosynthesis of previtamin D<sub>3</sub> in the skin. Atmospheric transmission of UVB is strongly influenced by atmospheric conditions and solar zenith angle. We investigated the effects of diurnal and seasonal variation on the availability of sufficient UVB radiation for adequate previtamin D<sub>3</sub> synthesis using an established in vitro model. Borosilicate ampoules of 7-dehydrocholesterol, the precursor of previtamin D<sub>3</sub>, in ethanol (50 µg/mL) were exposed to direct sunlight in an urban area of Abu Dhabi, at one hourly intervals between 0800 and 1700, on one day of each month over a period of one year. Conversion to previtamin D<sub>3</sub>, vitamin D<sub>3</sub> and metabolically inactive photoisomers was analyzed using high performance liquid chromatography. The efficiency of 7-dehydrocholesterol conversion to previtamin D<sub>3</sub> varied estimated UVB intensity. At the latitude of Abu Dhabi (24.2°N) previtamin D<sub>3</sub> synthesis can occur throughout the year. However very little if any previtamin D<sub>3</sub> was produced before 0900 hrs. and after 1600 hrs. Local conditions in Abu Dhabi are likely sufficient to maintain vitamin D levels throughout the year given adequate sun exposure.

### ARTICLE HISTORY

Received 27 October 2016  
Accepted 28 November 2016

### KEYWORDS

7-dehydrocholesterol;  
atmospheric modeling;  
lumisterol; previtamin D<sub>3</sub>;  
tachysterol; UVB; vitamin D<sub>3</sub>

### Introduction

Despite the increasing public awareness of the potential health benefits of adequate vitamin D intake, vitamin D deficiency remains endemic worldwide.<sup>1</sup> The active form, 1,25-dihydroxyvitamin D (1,25(OH)<sub>2</sub>D), plays a crucial role in metabolic processes ranging from calcium and phosphorous metabolism to cell maturation and growth.

Given sufficient sunlight containing UVB radiation the 7-dehydrocholesterol in the skin is converted to previtamin D.<sup>2</sup> 7-dehydrocholesterol is isomerised to previtamin D<sub>3</sub> by photons at wavelengths of 290–315nm, corresponding to the ultraviolet B (UVB) range of the electromagnetic spectrum. A heat-dependent process converts previtamin D<sub>3</sub> (which is thermodynamically unstable) to vitamin D<sub>3</sub>, while further exposure to UVB radiation photoisomerises previtamin D<sub>3</sub> into lumisterol<sub>3</sub>, tachysterol<sub>3</sub> or 7-dehydrocholesterol. Consequently, even brief exposure to sunlight allows for the synthesis of vitamin D<sub>3</sub>, while protracted

sunlight exposure leads to production of the biologically inert isomers, averting vitamin D toxicity.<sup>3</sup> Vitamin D<sub>3</sub> is itself degraded by solar irradiation to the metabolically inactive suprasterol 1 and 2,<sup>4</sup> providing an additional safeguard against toxicity in conditions of high sunlight exposure. Vitamin D<sub>3</sub>, together with vitamin D<sub>2</sub> acquired from dietary sources, undergoes 25-hydroxylation in the liver by Cyp2R1 to 25-hydroxyvitamin D (25(OH)D). 25(OH)D is the major circulatory form of vitamin D and correlates with total body stores, and hence is used as the substrate for most commercially available vitamin D assays.<sup>5</sup> The metabolically active form of vitamin D, 1,25-dihydroxyvitamin D (1,25(OH)<sub>2</sub>D), is synthesized from 25(OH)D by further hydroxylation in the kidney.

Dietary vitamin D intake is usually insufficient to meet requirements and therefore most humans remain dependent on photosynthesis.<sup>6</sup> UVB intensity at ground level is determined predominantly by solar zenith angle and atmospheric ozone. In general, terrain has a low albedo

**CONTACT** Nader Lessan, MD  [nlessan@icldc.ae](mailto:nlessan@icldc.ae)  FRCP, Consultant Endocrinologist, Imperial College London Diabetes Centre, PO Box 48338, Abu Dhabi, United Arab Emirates.

© 2018 Adam J. Buckley, Zara Hannoun, Nader Lessan, Michael F. Holick, and Maha T. Barakat. Published with license by Taylor & Francis.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

to UV and hence reflected rays do not contribute significantly to local UVB intensity.<sup>7</sup> While shade from trees permits sufficient transmission of UVB for vitamin D synthesis, shade from buildings and glass windows significantly attenuates biologically significant wavelengths.<sup>8</sup> Cultural differences in clothing, and consequently the area of skin exposed to sunlight, affect vitamin D production when overall UVB exposure is low, although the effect of clothing is greatly reduced when there is an excess of available UVB.<sup>9</sup> Melanin absorbs electromagnetic radiation at wavelengths of 280–400 nm, and hence darker skin pigmentation can diminish the potential for vitamin D synthesis.<sup>10</sup> Assessment of individual vitamin D status is complicated by polymorphisms of the vitamin D binding protein,<sup>11,12</sup> and by sequestration of the fat-soluble vitamin D in adipose tissue.<sup>13</sup>

Vitamin D deficiency is highly prevalent in the UAE population. A limited number of studies have investigated the prevalence of, and local factors associated with, vitamin D deficiency. A study conducted in Al Ain demonstrated hypovitaminosis D in 65.1% of a population of 143 adolescents, with vitamin D deficiency  $\leq 50$  nmol/l in 10% of boys and 28% of girls.<sup>14</sup> Vitamin D status was found to be paradoxically lower in summer than in winter in a population of 138 female university students in Abu Dhabi, in whom sun avoidance behavior was common and independently negatively correlated with vitamin D status.<sup>15</sup> Of children aged 8–14 attending an ambulatory clinic at Sheikh Khalifa Medical City in Abu Dhabi, 31.2% had evidence of severe vitamin D deficiency with serum 25(OH)D  $< 25$  nmol/l.<sup>16</sup>

We hypothesized that local environmental and atmospheric factors such as dust and ground level ozone might contribute to the high local prevalence of vitamin D deficiency; we explored this hypothesis by examining seasonal and diurnal variation in photosynthesis of vitamin D in an established *in vitro* model, and compared the results with predicted conversion of 7-dehydrocholesterol to previtamin D<sub>3</sub> based on atmospheric modeling and the action spectrum of UVB for photoisomerisation of 7-dehydrocholesterol to previtamin D<sub>3</sub>.

## Materials and methods

### *In vitro* conversion of 7-dehydrocholesterol

Borosilicate glass ampoules of 7-dehydrocholesterol dissolved in ethanol (50  $\mu$ g/mL) were sealed under

argon as previously described,<sup>17</sup> and stored in a cool dark location prior to use. Each ampoule was placed outside in direct sunlight, for a period of one hour, at hourly intervals between 8:00 to 16:00 on the last Wednesday of each month of 2012. Once exposed, ampoules were stored in darkness at  $-4^{\circ}\text{C}$  pending analysis. The proportions of 7-dehydrocholesterol, previtamin D and vitamin D, tachysterol and lumisterol in each sample were assayed using high performance liquid chromatography (Boston University Hospital, Boston, USA) as previously described.<sup>18</sup>

### *Atmospheric model and predicted 7-dehydrocholesterol conversion*

We developed a model to predict the potential for incident UVB to convert 7-dehydrocholesterol into previtamin D<sub>3</sub> at each of the time points sampled. Sufficiently detailed meteorological data during the period of study were not available, and it was therefore necessary to estimate UVB intensity. A simplified model of clear-sky spectral irradiance at the earth's surface, SPCTRL2,<sup>19</sup> as implemented in the C programming language by M. Rimes (NREL, 1998),<sup>20</sup> was used to estimate irradiance of a horizontal surface at the latitude and longitude of Abu Dhabi (24.4667° N, 54.3667° E). Total precipitable water content was estimated at 1.5 cm. The SPCTRL2 default setting for atmospheric ozone was used. Surface albedo was estimated at 0.1. Spectral irradiance was modeled between 290 and 310 nm; at wavelengths shorter than 300 nm very little UV penetrates to ground level due to absorption by atmospheric ozone,<sup>21</sup> while wavelengths longer than 310 nm convert 7-dehydrocholesterol to vitamin D<sub>3</sub> with very low efficiency.<sup>22,23</sup> SPCTRL2 estimates of combined direct and diffuse incident energy at wavelengths of 290, 295, 300, 305 and 310 nm were obtained for each studied time period, expressed in  $\text{Js}^{-1}\text{m}^2\text{nm}^{-1}$ , and linearly interpolated to determine the number of joules of energy delivered at each 1 nm interval of wavelength over one hour. These estimates were weighted by the action spectrum of UVB on 7-dehydrocholesterol,<sup>23</sup> and interpolated with the assumption of a quadratic relationship between required incident energy and conversion at wavelengths between 300 and 310 nm. The results were summed, integrating the potential for photoisomerisation of 7-dehydrocholesterol by the 290–310 nm spectrum and providing an estimate of total percentage conversion to previtamin D during each time interval studied.

## Results

### Diurnal variation


The conversion of 7-dehydrocholesterol to previtamin D varied with time of day on each individual day studied. Over the 12 sampled days, conversion was negligible during the 0800–0900 (mean 0.53%, range 0.0–0.94%) and 1600–1700 (mean 0.33%, range 0.0–1.0%) time periods, and peaked during the 1200–1300 time period (mean 4.8%, range 4.7–11.0%). Rates of conversion to tachysterol and lumisterol followed a similar pattern, although absolute values were smaller. The conversion followed a sinusoidal pattern on each day studied, indicating that zenith angle was the most important determinant (see Fig. 1).

By extrapolation we predicted that very little conversion of 7-dehydrocholesterol would occur outside the studied time periods. Variation in the proportion of 7-dehydrocholesterol converted during each month can therefore be expressed in terms of the area under


the curve of the graphs of diurnal variation in rate of conversion. Total daily conversion to previtamin D and 25(OH)D, expressed as the area under the curve of conversion at each time point studied, was 2.92 times higher in July than in January (see Fig. 2).

### Observed vs. predicted conversion of 7-dehydrocholesterol to previtamin D3

Linear regression of predicted against actual conversion rates showed a strong and highly significant association ( $r^2 = 0.82$ ,  $p < 0.0001$ , see Fig. 3). On average the model underestimated conversion, (predicted =  $0.79 \times \text{observed} + 0.12$ ). This may be explained by the assumption of low surface albedo, excluding the contribution of reflected light, which, while true for a natural environment, may not apply equally to urban conditions. Linear regression of predicted against actual conversion for each time period indicated that the model tended to underestimate conversion during




**Figure 1.** Diurnal variation in 7-dehydrocholesterol conversion in vitro to previtamin D3 or vitamin D3, lumisterol or tachysterol (%) following 1 hour exposure to sunlight, during each month of the year.



**Figure 2.** Seasonal variation in in vitro 7-dehydrocholesterol conversion to previtamin D, expressed as area under the curve of percentage conversion throughout the day in each month. Values are normalized to July.

the morning and overestimate conversion during the evening. Incident UVB at a point on the Earth's surface varies with time of day according to a sinusoidal, rather than linear pattern and hence this model would tend to underestimate while incident UVB was increasing and overestimate while incident UVB was decreasing. This model made a number of simplifying




**Figure 3.** Observed conversion of 7-dehydrocholesterol to previtamin D3 and vitamin D3 (%) plotted against conversion predicted by the study model, based on estimated irradiance and action spectrum of UVB on conversion (%). Predicted =  $(0.794 \times \text{observed}) + 0.123$ ,  $R^2 = 0.823$ ,  $p < 0.0001$ .

assumptions, including a quadratic relationship between UVB intensity and rate of conversion and linear interpolation of UV intensities, and therefore while the significant correlation between the predicted and observed values indicates that incident UV is likely to be the most important factor influencing conversion, the remarkable numerical similarity of the predicted and observed values is likely to be coincidental.

## Discussion

The conversion of 7-dehydrocholesterol to previtamin D3 and its photoproducts in vitro seen in Abu Dhabi are comparable with similar ampoule studies performed in the Middle East and around the world, in locations including Boston, Johannesburg, Riyadh, South India and Argentina (see Fig. 4).<sup>17,24-27</sup> It has previously been established that the results of the ampoule model correlate with those observed in human skin samples exposed under the same conditions.<sup>17,28</sup>

Conversion of 7-dehydrocholesterol to previtamin D in vitro was well correlated with local availability of radiation in the UVB spectrum, and the percentage of previtamin D produced in each ampoule was equal to or exceeded that predicted by our model for each corresponding time period. Although atmospheric ozone was estimated, it would be expected that significant ground level ozone would cause measured conversion to be less, rather than greater than, predicted values. This pattern of variation suggests that local environmental factors, such as dust and ground ozone, do not attenuate incident UVB to an extent sufficient to limit previtamin D synthesis. In addition, the conversion of 7-dehydrocholesterol to previtamin D3 seen in Abu Dhabi are comparable with other sunny locations around the world, where vitamin D deficiency is



**Figure 4.** Seasonal variation in 7-dehydrocholesterol conversion in vitro in ampoules exposed to direct sunlight between 12:00 and 13:00 worldwide. Data supplied by MF Holick.

considerably rarer.<sup>29-31</sup> Other explanations for the high prevalence of vitamin D deficiency seen in the UAE therefore need to be considered.

## Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

## Funding

The study was funded by Mubadala Healthcare.

## Author contributions

AB wrote the first draft of the manuscript and analyzed the data. ZH performed the in vitro study and contributed to the manuscript. NL reviewed and revised the manuscript. MH supplied the study ampoules and performed the assays, and read and revised the manuscript. MB reviewed and commented on the manuscript. AB had full access to study data and as such is guarantor of the work.

## ORCID

Adam J. Buckley  <http://orcid.org/0000-0001-7471-0273>

Michael F. Holick  <http://orcid.org/0000-0001-6023-9062>

## References

- [1] Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutri* 2008; 87:1080S; PMID:18400738
- [2] Glossmann HH. Origin of 7-Dehydrocholesterol (Previtamin D) in the Skin. *J Invest Dermatol* 2010; 130:2139-41; PMID:20445550; <https://doi.org/10.1038/jid.2010.118>
- [3] Holick MF. The cutaneous photosynthesis of previtamin D3: a unique photoendocrine system. *J Investigative Dermatol* 1981; 77:51-8; PMID:6265564; <https://doi.org/10.1111/1523-1747.ep12479237>
- [4] Webb AR, DeCosta BR, Holick MF. Sunlight regulates the cutaneous production of vitamin D3 by causing its photo-degradation. *J Clin Endocrinol Metab* 1989; 68:882-7; PMID:2541158; <https://doi.org/10.1210/jcem-68-5-882>
- [5] Carter GD. Accuracy of 25-hydroxyvitamin D assays: confronting the issues. *Curr Drug Targets* 2011; 12:19-28; PMID:20795940; <https://doi.org/10.2174/138945011793591608>
- [6] Hollis BW. Circulating 25-hydroxyvitamin D levels indicative of vitamin D sufficiency: implications for establishing a new effective dietary intake recommendation for vitamin D. *J Nutri* 2005; 135:317-22; PMID:15671234
- [7] Allaart M, van Weele M, Fortuin P, Kelder H. An empirical model to predict the UV-index based on solar zenith angles and total ozone. *Meteorological Applications* 2004; 11:59-65; <https://doi.org/10.1017/S1350482703001130>
- [8] Turnbull DJ, Parisi AV, Kimlin MG. Vitamin D effective ultraviolet wavelengths due to scattering in shade. *J Steroid Biochem Mol Biol* 2005; 96:431-6; PMID:16005208; <https://doi.org/10.1016/j.jsbmb.2005.04.039>
- [9] Bogh MKB, Schmedes AV, Philipsen PA, Thieden E, Wulf HC. Interdependence between body surface area and ultraviolet B dose in vitamin D production: a randomized controlled trial. *Br J of Dermatol* 2010; 164:163-9; <https://doi.org/10.1111/j.1365-2133.2010.10082.x>
- [10] Dawson-Hughes B. Serum 25-hydroxyvitamin D and functional outcomes in the elderly. *Am J Clin Nutri* 2008; 88:537-40
- [11] Perna L, Felix JF, Breitling LP, Haug U, Raum E, Burwinkel B, Schöttker B, Brenner H. Genetic variations in the vitamin D binding protein and season-specific levels of vitamin D among older adults. *Epidemiol (Cambridge, Mass)* 2013; 24:104-9; PMID:23191998; <https://doi.org/10.1097/EDE.0b013e318276c4b0>
- [12] Powe CE, Evans MK, Wenger J, Zonderman AB, Berg AH, Nalls M, Tamez H, Zhang D, Bhan I, Karumanchi SA, et al. Vitamin D-Binding Protein and Vitamin D Status of Black Americans and White Americans. *N Eng J Med* 2013; 369:1991-2000; PMID:24256378; <https://doi.org/10.1056/NEJMoa1306357>
- [13] Holick MF. Vitamin D: importance in the prevention of cancers, type 1 diabetes, heart disease, and osteoporosis. *Am J Clin Nutri* 2004; 79:362-71; PMID:14985208
- [14] Muhairi SJ, Mehairi AE, Khouri AA, Naqbi MM, Maskari FA, Al Kaabi J, Al Dhaheri AS, Nagelkerke N, Shah SM. Vitamin D deficiency among healthy adolescents in Al Ain, United Arab Emirates. *BMC Public Health* 2013; 13:33; PMID:23311702; <https://doi.org/10.1186/1471-2458-13-33>
- [15] Anouti F Al, Thomas J, Abdel-Wareth L, Rajah J, Grant WB, Haq A. Vitamin D deficiency and sun avoidance among university students at Abu Dhabi, United Arab Emirates. *Dermato-Endocrinol* 2011; 3:235-9; PMID:22259650; <https://doi.org/10.4161/derm.3.4.16881>
- [16] Rajah J, Haq A, Pettifor JM. Vitamin D and calcium status in urban children attending an ambulatory clinic service in the United Arab Emirates. *Dermato-Endocrinol* 2012; 4:39-43; PMID:22870351; <https://doi.org/10.4161/derm.18250>
- [17] Webb AR, Kline L, Holick MF. Influence of season and latitude on the cutaneous synthesis of vitamin D3: exposure to winter sunlight in Boston and Edmonton will not promote vitamin D3 synthesis in human skin. *J Clin Endocrinol Metab* 1988; 67:373-8; PMID:2839537; <https://doi.org/10.1210/jcem-67-2-373>
- [18] Holick M, MacLaughlin J, Clark M, Holick SA, Potts JT, Jr, Anderson RR, Blank IH, Parrish JA, Elias P. Photosynthesis of previtamin D3 in human skin and the physiologic consequences. *Science* 1980; 210:203-5; PMID:6251551; <https://doi.org/10.1126/science.6251551>
- [19] Bird RE, Riordan C. Simple solar spectral model for direct and diffuse irradiance on horizontal and tilted planes at the earth's surface for cloudless atmospheres. *J Climate Applied Meteo* 1986; 25(1):87-97.

- [20] Rymes M, Wilcox S, Anderberg M. SPCTRL2. 1998; Golden (CO): National Renewable Energy Laboratory. Available from: <http://rredc.nrel.gov/solar/models/spectral/spectrl2/>
- [21] Leckner B. The spectral distribution of solar radiation at the earth's surface—elements of a model. *Solar Energy* 1978; 20(2):143-50.
- [22] Wacker M, Holick MF. Sunlight and Vitamin D: A global perspective for health. *Dermato-Endocrinol* 2013; 5:51-108; PMID:24494042; <https://doi.org/10.4161/derm.24494>
- [23] Bolsée D, Webb AR, Gillotay D, Dörschel B, Knuschke P, Krins A, Terenetskaya I. Laboratory facilities and recommendations for the characterization of biological ultraviolet dosimeters. *Applied Optics* 2000; 39:2813-22; <https://doi.org/10.1364/AO.39.002813>
- [24] Harinarayan CV, Holick MF, Prasad UV, Vani PS, Himabindu G. Vitamin D status and sun exposure in India. *Dermato-Endocrinol* 2013; 5:130-41; PMID:24494046; <https://doi.org/10.4161/derm.23873>
- [25] Pettifor JM, Moodley GP, Hough FS, Koch H, Chen T, Lu Z, Holick MF. The effect of season and latitude on in vitro vitamin D formation by sunlight in South Africa. *South African Medical J = Suid-Afrikaanse tydskrif vir geneeskunde* 1996; 86:1270-2; PMID:8955733
- [26] Ladizesky M, Lu Z, Oliveri B, San Roman N, Diaz S, Holick MF, Mautalen C. Solar ultraviolet B radiation and photoproduction of vitamin D3 in central and southern areas of argentina. *J Bone Mineral Res* 1995; 10:545-9; PMID:7610924; <https://doi.org/10.1002/jbmr.5650100406>
- [27] Alshahrani FM, Almalki MH, Aljohani N, Alzahrani A, Alsaleh Y, Holick MF. Vitamin D: Light side and best time of sunshine in Riyadh, Saudi Arabia. *Dermato-Endocrinology* 2013; 5:177-80; PMID:24494051; <https://doi.org/10.4161/derm.23351>
- [28] Chen TC, Lu Z, Holick MF. Photobiology of Vitamin D. *Vitamin D* 2010:35-60; [https://doi.org/10.1007/978-1-60327-303-9\\_2](https://doi.org/10.1007/978-1-60327-303-9_2)
- [29] Levis S, Gomez A, Jimenez C, Veras L, Ma F, Lai S, Hollis B, Roos BA. Vitamin D Deficiency and Seasonal Variation in an Adult South Florida Population. *J Clin Endocrinol Metab* 2005; 90:1557-62; PMID:15634725; <https://doi.org/10.1210/jc.2004-0746>
- [30] Poopedi MA, Norris SA, Pettifor JM. Factors influencing the vitamin D status of 10-year-old urban South African children. *Public Health Nutrition* 2011; 14:334-9; PMID:20804632; <https://doi.org/10.1017/S136898001000234X>
- [31] Fradinger EE, Zanchetta JR. Vitamin D and bone mineral density in ambulatory women living in Buenos Aires, Argentina. *Osteoporosis Int* 2001; 12:24-7; PMID:11305079; <https://doi.org/10.1007/s001980170153>